arXiv:q-alg/9703031v1 17 Mar 1997

Super Yangian Double $DY(gl(1|1))$ With Center Extension

[a]Jin-fang Cai, [a]Guo-xing Ju [bc]S.K. Wang and [a]Ke Wu

[a] Institute of Theoretical Physics, Academia Sinica,
Beijing, 100080, P. R. China

[b] CCAST (World Laboratory), P.O. Box 3730, Beijing, 100080, P. R. China

[c] Institute of Applied Mathematics, Academia Sinica,
Beijing, 100080, P. R. China

Abstract

We use the super version of RS construction method and Gauss decomposition to obtain Drinfel′d's currents realization of super Yangian double $DY(gl(1|1))$ with center extension

Yangians and quantum affine algebras are quantum algebras which are related respectively with rational and trigonometric solutions of Yang-Baxter equation [1]. The explicit isomorphism between two realizations of quantum affine algebras and Yangians given by Drinfel′d [2] and RS [3] was established by Ding and Frenkel [4] using Gauss decomposition. In this paper, we use super version of RS construction method to define super Yangian double $DY(gl(1|1))$ [5] with center extension, and get its Drinfeld's currents realization by Guass decomposition.

By V we denote a $\mathbf{Z}_2$-graded two-dimension vector space (graded auxiliary space), and set the first and second basis of V is even and odd respectively. In graded case, we must use this tensor product form: $(A \otimes B)(C \otimes D) = (-1)^{P(B)P(C)} AC \otimes BD$, here $P(A) = 0, 1$ when A is bosonic and Fermionic respectively. Then the graded(super) Yang-Baxter equation(YBE) takes this form[6]:

$$\eta_{12}R_{12}(u)\eta_{13}R_{13}(u+v)\eta_{23}R_{23}(v) = \eta_{23}R_{23}(v)\eta_{13}R_{13}(u+v)\eta_{12}R_{12}(u), \tag{1}$$

where $R(u) \in End(V \otimes V)$ and it must obey weight conservation condition: $R_{ij,kl} \neq 0$ only when $i + j = k + l$. $\eta_{ik,jl} = (-1)^{(i-1)(k-1)}\delta_{ij}\delta_{lk}$. It's easily proved that the following $R(u)$ matrix is a rational solution of super YBE:

$$R(u) = \frac{1}{u+\hbar}(uI + \mathcal{P}) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \frac{u}{u+\hbar} & \frac{\hbar}{u+\hbar} & 0 \\ 0 & \frac{\hbar}{u+\hbar} & \frac{u}{u+\hbar} & 0 \\ 0 & 0 & 0 & \frac{u-\hbar}{u+\hbar} \end{pmatrix} \tag{2}$$

This sulution of graded YBE satisfied unitarity condition: $R(u)R_{21}(-u) = I$. From this rational solution of graded YBE, we can define the super Yangian double $DY(gl(1|1))$ with a central extension employing super version of RS method [3]. Super Yangian Double $DY\left(gl(1|1)\right)$ is a Hopf algebra generated by $\{l_{ij}^k | 1 \leq i,j \leq 2, k \in \mathbf{Z}\}$ which obey the following relations:

$$R(u-v)L_1^{\pm}(u)\eta L_2^{\pm}(v)\eta = \eta L_2^{\pm}(v)\eta L_1^{\pm}(u)R(u-v) \tag{3}$$

$$R(u_- - v_+)L_1^{+}(u)\eta L_2^{-}(v)\eta = \eta L_2^{-}(v)\eta L_1^{+}(u)R(u_+ - v_-) \tag{4}$$

here $u_\pm = u \pm \frac{\hbar}{4}c$ and we have used standard notation: $L_1^{\pm}(u) = L^{\pm}(u) \otimes \mathbf{1}, L_2^{\pm}(u) = \mathbf{1} \otimes L^{\pm}(u)$ and $L^{\pm}(u) = \left(l_{ij}^{\pm}(u)\right)_{1\leq i,j\leq 2}$, $l_{ij}^{\pm}(u)$ are generate functions of l_{ij}^k :

$$l_{ij}^{+}(u) = \delta_{ij} - \hbar \sum_{k\geq 0} l_{ij}^k u^{-k-1}, \quad l_{ij}^{-}(u) = \delta_{ij} + \hbar \sum_{k<0} l_{ij}^k u^{-k-1}. \tag{5}$$

Hopf structure of $DY\left(gl(1|1)\right)$ are defined as follows:

$$\triangle\left(l_{ij}^{\pm}(u)\right) = \sum_{k=1,2} l_{kj}^{\pm}(u \pm \frac{\hbar}{4}c_2) \otimes l_{ik}^{\pm}(u \mp \frac{\hbar}{4}c_1)(-1)^{(k+i)(k+j)}, \tag{6}$$

$$\epsilon\left(l_{ij}^{\pm}(u)\right) = \delta_{ij}, \qquad S\left({}^{st}L^{\pm}(u)\right) = \left[{}^{st}L^{\pm}(u)\right]^{-1}. \tag{7}$$

here $\left[{}^{st}L^{\pm}(u)\right]_{ij} = (-1)^{i+j} l_{ji}^{\pm}(u)$.

From Ding-Frenkel theorem[4] on two realizations of Yangians and quantum affine algebras, $L^{\pm}(u)$ have the following decomposition:

$$\begin{aligned} L^{\pm}(u) &= \begin{pmatrix} 1 & 0 \\ f^{\pm}(u) & 1 \end{pmatrix} \begin{pmatrix} k_1^{\pm}(u) & 0 \\ 0 & k_2^{\pm}(u) \end{pmatrix} \begin{pmatrix} 1 & e^{\pm}(u) \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} k_1^{\pm}(u) & k_1^{\pm}(u)e^{\pm}(u) \\ f^{\pm}(u)k_1^{\pm}(u) & k_2^{\pm}(u) + f^{\pm}(u)k_1^{\pm}(u)e^{\pm}(u) \end{pmatrix} \end{aligned} \tag{8}$$

and the inversions of $L^{\pm}(u)$ can be writed as:

$$\begin{aligned} L^{\pm}(u)^{-1} &= \begin{pmatrix} 1 & -e^{\pm}(u) \\ 0 & 1 \end{pmatrix} \begin{pmatrix} k_1^{\pm}(u)^{-1} & 0 \\ 0 & k_2^{\pm}(u)^{-1} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -f^{\pm}(u) & 1 \end{pmatrix} \\ &= \begin{pmatrix} k_1^{\pm}(u)^{-1} + e^{\pm}(u)k_2^{\pm}(u)^{-1}f^{\pm}(u) & -e^{\pm}(u)k_2^{\pm}(u)^{-1} \\ -k_2^{\pm}(u)^{-1}f^{\pm}(u) & k_2^{\pm}(u)^{-1} \end{pmatrix} \end{aligned} \tag{9}$$

Let

$$X^{+}(v) = e^{+}(v_-) - e^{-}(v_+) \qquad X^{-}(v) = f^{+}(v_+) - f^{-}(v_-) \tag{10}$$

To calculate the (anti-)commutation relations of $X^{\pm}(u)$ and $k_i^{\pm}(v)$ (i=1,2), we will use the following equivalent forms of eq.(3)(4):

$$L_1^{\pm}(u)^{-1}\eta L_2^{\pm}(v)^{-1}\eta R(u-v) = R(u-v)\eta L_2^{\pm}(v)^{-1}\eta L_1^{\pm}(u)^{-1} \tag{11}$$

$$L_1^{+}(u)^{-1}\eta L_2^{-}(v)^{-1}\eta R(u_- - v_+) = R(u_+ - v_-)\eta L_2^{-}(v)^{-1}\eta L_1^{+}(u)^{-1} \tag{12}$$

$$L_1^{\pm}(u)R(u-v)\eta L_2^{\pm}(v)^{-1}\eta = \eta L_2^{\pm}(v)^{-1}\eta R(u-v)L_1^{\pm}(u) \tag{13}$$

$$L_1^{+}(u)R(u_+ - v_-)\eta L_2^{-}(v)^{-1}\eta = \eta L_2^{-}(v)^{-1}\eta R(u_- - v_+)L_1^{+}(u) \tag{14}$$

$$L_1^{\pm}(u)^{-1}R_{21}(v-u)\eta L_2^{\pm}(v)\eta = \eta L_2^{\pm}(v)\eta R_{21}(v-u)L_1^{\pm}(u)^{-1} \tag{15}$$

$$L_1^{+}(u)^{-1}R_{21}(v_+ - u_-)\eta L_2^{-}(v)\eta = \eta L_2^{-}(v)\eta R_{21}(v_- - u_+)L_1^{+}(u)^{-1} \tag{16}$$

From (3) (4) and (11)—(16), we can get all relations between $k_1^{\pm}(u)$ and $k_2^{\pm}(v)$:

$$[k_1^{\pm}(u)\ ,\ k_1^{\pm}(v)] = [k_1^{+}(u)\ ,\ k_1^{-}(v)] = 0 \tag{17}$$

$$[k_1^{\pm}(u)\ ,\ k_2^{\pm}(v)] = [k_2^{\pm}(u)\ ,\ k_2^{\pm}(v)] = 0 \tag{18}$$

$$\frac{u_{\pm} - v_{\mp}}{u_{\pm} - v_{\mp} + \hbar} k_1^{\pm}(u) k_2^{\mp}(v)^{-1} = \frac{u_{\mp} - v_{\pm}}{u_{\mp} - v_{\pm} + \hbar} k_2^{\mp}(v)^{-1} k_1^{\pm}(u) \tag{19}$$

$$\frac{u_{-} - v_{+} - \hbar}{u_{-} - v_{+} + \hbar} k_2^{+}(u)^{-1} k_2^{-}(v)^{-1} = \frac{u_{+} - v_{-} - \hbar}{u_{+} - v_{-} + \hbar} k_2^{-}(v)^{-1} k_2^{+}(u)^{-1} \tag{20}$$

From (3)(4) and unitarity of R-matrix, we have the following relations between $k_1^{\pm}(u)$ and $e^{\pm}(v), f^{\pm}(v)$:

$$\begin{aligned} k_1^{\pm}(u) k_1^{\pm}(v) e^{\pm}(v) - \frac{u - v}{u - v + \hbar} k_1^{\pm}(v) e^{\pm}(v) k_1^{\pm}(u) \\ - \frac{\hbar}{u - v + \hbar} k_1^{\pm}(v) k_1^{\pm}(u) e^{\pm}(u) = 0 \end{aligned} \tag{21}$$

$$\begin{aligned} k_1^{\pm}(u) k_1^{\mp}(v) e^{\mp}(v) - \frac{u_{\pm} - v_{\mp}}{u_{\pm} - v_{\mp} + \hbar} k_1^{\mp}(v) e^{\mp}(v) k_1^{\pm}(u) \\ - \frac{\hbar}{u_{\pm} - v_{\mp} + \hbar} k_1^{\mp}(v) k_1^{\pm}(u) e^{\pm}(u) = 0 \end{aligned} \tag{22}$$

$$\begin{aligned} f^{\pm}(v) k_1^{\pm}(v) k_1^{\pm}(u) - \frac{u - v}{u - v + \hbar} k_1(u) f^{\pm}(v) k_1^{\pm}(v) \\ - \frac{\hbar}{u - v + \hbar} f^{\pm}(u) k_1^{\pm}(u) k_1^{\pm}(v) = 0 \end{aligned} \tag{23}$$

$$\begin{aligned} f^{\mp}(v) k_1^{\mp}(v) k_1^{\pm}(u) - \frac{u_{\mp} - v_{\pm}}{u_{\mp} - v_{\pm} + \hbar} k_1^{\pm}(u) f^{\mp}(v) k_1^{\mp}(v) \\ - \frac{\hbar}{u_{\mp} - v_{\pm} + \hbar} f^{\pm}(u) k_1^{\pm}(u) k_1^{\mp}(v) = 0 \end{aligned} \tag{24}$$

thus

$$(u_{\pm} - v + \hbar) e^{\pm}(v_{\mp}) - (u_{\pm} - v) k_1^{\pm}(u)^{-1} e^{\pm}(v_{\mp}) k_1^{\pm}(u) - \hbar e^{\pm}(u) = 0 \tag{25}$$

$$(u_{\pm} - v + \hbar) e^{\mp}(v_{\pm}) - (u_{\pm} - v) k_1^{\pm}(u)^{-1} e^{\mp}(v_{\pm}) k_1^{\pm}(u) - \hbar e^{\pm}(u) = 0 \tag{26}$$

$$(u_{\mp} - v + \hbar) f^{\pm}(v_{\pm}) - (u_{\mp} - v) k_1^{\pm}(u) f^{\pm}(v_{\pm}) k_1^{\pm}(u)^{-1} - \hbar f^{\pm}(u) = 0 \tag{27}$$

$$(u_{\mp} - v + \hbar) f^{\mp}(v_{\mp}) - (u_{\mp} - v) k_1^{\pm}(u) f^{\mp}(v_{\mp}) k_1^{\pm}(u)^{-1} - \hbar f^{\pm}(u) = 0 \tag{28}$$

so

$$k_1^{\pm}(u)^{-1} X^{+}(v) k_1 \pm (u) = \frac{u_{\pm} - v + \hbar}{u_{\pm} - v} X^{+}(v) \tag{29}$$

$$k_1^{\pm}(u) X^{-}(v) k_1^{\pm}(u)^{-1} = \frac{u_{\mp} - v + \hbar}{u_{\mp} - v} X^{-}(v) \tag{30}$$

Now, we derive relations between $k_2^{\pm}(u)$ and $X^{\pm}(v)$. From(11)(12) and unitarity of R-matirx, we have

$$\frac{u - v - \hbar}{u - v + \hbar} e^{\pm}(u) k_2^{\pm}(u)^{-1} k_2^{\pm}(v)^{-1} - \frac{u - v}{u - v + \hbar} k_2^{\pm}(v)^{-1} e^{\pm}(u) k_2^{\pm}(u)^{-1}$$

$$+\frac{\hbar}{u-v+\hbar}e^{\pm}(v)k_2^{\pm}(v)^{-1}k_2^{\pm}(u)^{-1}=0 \tag{31}$$

$$\frac{u_{\mp}-v_{\pm}-\hbar}{u_{\mp}-v_{\pm}+\hbar}e^{\pm}(u)k_2^{\pm}(u)^{-1}k_2^{\mp}(v)^{-1}-\frac{u_{\pm}-v_{\mp}}{u_{\pm}-v_{\mp}+\hbar}k_2^{\mp}(v)^{-1}e^{\pm}(u)k_2^{\pm}(u)^{-1}$$

$$+\frac{\hbar}{u_{\pm}-v_{\mp}+\hbar}e^{\mp}(v)k_2^{\mp}(v)^{-1}k_2^{\pm}(u)^{-1}=0 \tag{32}$$

$$\frac{u-v-\hbar}{u-v+\hbar}k_2^{\pm}(v)^{-1}k_2^{\pm}(u)^{-1}f^{\pm}(u)-\frac{u-v}{u-v+\hbar}k_2^{\pm}(u)^{-1}f^{\pm}(u)k_2^{\pm}(v)^{-1}$$

$$+\frac{\hbar}{u-v+\hbar}k_2^{\pm}(u)^{-1}k_2^{\pm}(v)^{-1}f^{\pm}(v)=0 \tag{33}$$

$$\frac{u_{\pm}-v_{\mp}-\hbar}{u_{\pm}-v_{\mp}+\hbar}k_2^{\mp}(v)^{-1}k_2^{\pm}(u)^{-1}f^{\pm}(u)-\frac{u_{\mp}-v_{\pm}}{u_{\mp}-v_{\pm}+\hbar}k_2^{\pm}(u)^{-1}f^{\pm}(u)k_2^{\mp}(v)^{-1}$$

$$+\frac{\hbar}{u_{\mp}-v_{\pm}+\hbar}k_2^{\pm}(u)^{-1}k_2^{\mp}(v)^{-1}f^{\mp}(v)=0 \tag{34}$$

Using relations between $k_2^{+}(u)$ and $k_2^{-}(v)$ (18)(20), we have

$$(u_{\mp}-v-\hbar)e^{\pm}(u_{\mp})-(u_{\mp}-v)k_2^{\pm}(v)^{-1}e^{\pm}(u_{\mp})k_2^{\pm}(v)+\hbar e^{\pm}(v)=0 \tag{35}$$

$$(u-v_{\mp}-\hbar)e^{\pm}(u_{\mp})-(u-v_{\mp})k_2^{\mp}(v)^{-1}e^{\pm}(u_{\mp})k_2^{\mp}(v)+\hbar e^{\mp}(v)=0 \tag{36}$$

$$(u_{\pm}-v-\hbar)f^{\pm}(u_{\pm})-(u_{\pm}-v)k_2^{\pm}(v)f^{\pm}(u_{\pm})k_2^{\pm}(v)^{-1}+\hbar f^{\pm}(v)=0 \tag{37}$$

$$(u-v_{\pm}-\hbar)f^{\pm}(u_{\pm})-(u-v_{\pm})k_2^{\mp}(v)f^{\pm}(u_{\pm})k_2^{\mp}(v)^{-1}+\hbar f^{\mp}(v)=0 \tag{38}$$

then

$$k_2^{\pm}(u)^{-1}X^{+}(v)k_2^{\pm}(u)=\frac{u_{\pm}-v+\hbar}{u_{\pm}-v}X^{+}(v) \tag{39}$$

$$k_2^{\pm}(u)X^{-}(v)k_2^{\pm}(u)^{-1}=\frac{u_{\mp}-v+\hbar}{u_{\mp}-v}X^{-}(v) \tag{40}$$

Now, we calculate the relations between $X^{\pm}(u)$ and $X^{\pm}(v)$. From (3)(4) and unitarity of R-matirx , we have

$$k_1^{\pm}(u)e^{\pm}(u)k_1^{\pm}(v)e^{\pm}(v)+\frac{u-v-\hbar}{u-v+\hbar}k_1^{\pm}(v)e^{\pm}(v)k_1^{\pm}(u)e^{\pm}(u)=0 \tag{41}$$

$$k_1^{\pm}(u)e^{\pm}(u)k_1^{\mp}(v)e^{\mp}(v)+\frac{u_{\pm}-v_{\mp}-\hbar}{u_{\pm}-v_{\mp}+\hbar}k_1^{\mp}(v)e^{\mp}(v)k_1^{\pm}(u)e^{\pm}(u)=0 \tag{42}$$

$$\frac{u-v-\hbar}{u-v+\hbar}f^{\pm}(u)k_1^{\pm}(u)f^{\pm}(v)k_1^{\pm}(v)+f^{\pm}(v)k_1^{\pm}(v)f^{\pm}(u)k_1^{\pm}(u)=0 \tag{43}$$

$$\frac{u_{\mp}-v_{\pm}-\hbar}{u_{\mp}-v_{\pm}+\hbar}f^{\pm}(u)k_1^{\pm}(u)f^{\mp}(v)k_1^{\mp}(v)+f^{\mp}(v)k_1^{\mp}(v)f^{\pm}(u)k_1^{\pm}(u)=0 \tag{44}$$

Using the above relations between $X^{\pm}(u)$ (or $e^{\pm}(u)$ and $f^{\pm}(u)$) and $k_1^{\pm}(v)$, it's easily to get the following relations;

$$X^{+}(u)X^{+}(v)+X^{+}(v)X^{+}(u)=0 \tag{45}$$

$$X^{-}(u)X^{-}(v)+X^{-}(v)X^{-}(u)=0 \tag{46}$$

From (3)(4) and unitarity of R-matirx, we have following relations :

$$\begin{aligned}&\frac{u-v}{u-v+\hbar}k_1^\pm(u)e^\pm(u)f^\pm(v)k_1^\pm(v)+\frac{u-v}{u-v+\hbar}f^\pm(v)k_1^\pm(v)k_1^\pm(u)e^\pm(u)\\&\qquad-\frac{\hbar}{u-v+\hbar}\left(k_2^\pm(u)+f^\pm(u)k_1^\pm(u)e^\pm(u)\right)k_1^\pm(v)\\&\qquad-\frac{\hbar}{u-v+\hbar}\left(k_2^\pm(v)+f^\pm(v)k_1^\pm(v)e^\pm(v)\right)k_1^\pm(u)=0\end{aligned}\tag{47}$$

$$\begin{aligned}&\frac{u_\mp-v_\pm}{u_\mp-v_\pm+\hbar}k_1^\pm(u)e^\pm(u)f^\mp(v)k_1^\mp(v)+\frac{u_\pm-v_\mp}{u_\pm-v_\mp+\hbar}f^\mp(v)k_1^\mp(v)k_1^\pm(u)e^\pm(u)\\&\qquad-\frac{\hbar}{u_\mp-v_\pm+\hbar}\left(k_2^\pm(u)+f^\pm(u)k_1^\pm(u)e^\pm(u)\right)k_1^\mp(v)\\&\qquad-\frac{\hbar}{u_\pm-v_\mp+\hbar}\left(k_2^\mp(v)+f^\mp(v)k_1^\mp(v)e^\mp(v)\right)k_1^\pm(u)=0\end{aligned}\tag{48}$$

then using (23)(24) and the following relations which also from (3)(4) and unitarity of R-matirx:

$$\begin{aligned}&k_1^\pm(u)e^\pm(u)k_1^\pm(v)-\frac{u-v}{u-v+\hbar}k_1^\pm(v)k_1^\pm(u)e^\pm(u)\\&\qquad-\frac{\hbar}{u-v+\hbar}k_1^\pm(v)e^\pm(v)k_1^\pm(u)=0\end{aligned}\tag{49}$$

$$\begin{aligned}&k_1^\pm(u)e^\pm(u)k_1^\mp(v)-\frac{u_\pm-v_\mp}{u_\pm-v_\mp+\hbar}k_1^\mp(v)k_1^\pm(u)e^\pm(u)\\&\qquad-\frac{\hbar}{u_\pm-v_\mp+\hbar}k_1^\mp(v)e^\mp(v)k_1^\pm(u)=0\end{aligned}\tag{50}$$

we obtain:

$$e^\pm(u)f^\pm(v)+f^\pm(v)e^\pm(u)=\frac{\hbar}{u-v}k_1^\pm(u)^{-1}k_2^\pm(u)-\frac{\hbar}{u-v}k_1^\pm(v)^{-1}k_2^\pm(v)\tag{51}$$

$$e^\pm(u)f^\mp(v)+f^\mp(v)e^\pm(u)=\frac{\hbar}{u_\mp-v_\pm}k_1^\pm(u)^{-1}k_2^\pm(u)-\frac{\hbar}{u_\pm-v_\mp}k_1^\mp(v)^{-1}k_2^\mp(v)\tag{52}$$

then

$$\{X^+(u)\ ,\ X^-(v)\}=\hbar\left[\delta(u_--v_+)k_1^+(u_-)^{-1}k_2^+(u_-)-\delta(u_+-v_-)k_1^-(v_-)^{-1}k_2^-(v_-)\right]\tag{53}$$

where $\delta(u-v)=\sum_{k\in Z}u^kv^{-k-1}$.

In a word, we get all relations between $k_i^\pm(u)$ (i=1,2) and $X^\pm(v)$:

$$[k_1^\pm(u)\ ,\ k_1^\pm(v)]=[k_1^+(u)\ ,\ k_1^-(v)]=0\tag{54}$$

$$[k_1^\pm(u)\ ,\ k_2^\pm(v)]=[k_2^\pm(u)\ ,\ k_2^\pm(v)]=0\tag{55}$$

$$\frac{u_\pm-v_\mp}{u_\pm-v_\mp+\hbar}k_1^\pm(u)k_2^\mp(v)^{-1}=\frac{u_\mp-v_\pm}{u_\mp-v_\pm+\hbar}k_2^\mp(v)^{-1}k_1^\pm(u)\tag{56}$$

$$\frac{u_--v_+-\hbar}{u_--v_++\hbar}k_2^+(u)^{-1}k_2^-(v)^{-1}=\frac{u_+-v_--\hbar}{u_+-v_-+\hbar}k_2^-(v)^{-1}k_2^+(u)^{-1}\tag{57}$$

$$k_i^\pm(u)^{-1}X^+(v)k_i^\pm(u)=\frac{u_\pm-v+\hbar}{u_\pm-v}X^+(v)\qquad(i=1,2)\tag{58}$$

$$k_i^{\pm}(u)X^-(v)k_i^{\pm}(u)^{-1} = \frac{u_{\mp} - v + \hbar}{u_{\mp} - v}X^-(v) \qquad (i = 1, 2) \tag{59}$$

$$\{X^+(u)\ ,\ X^+(v)\} = \{X^-(u)\ ,\ X^-(v)\} = 0 \tag{60}$$

$$\{X^+(u)\ ,\ X^-(v)\} = \hbar\left[\delta(u_- - v_+)k_1^+(u_-)^{-1}k_2^+(u_-) - \delta(u_+ - v_-)k_1^-(v_-)^{-1}k_2^-(v_-)\right] \tag{61}$$

Introducing a transformation for currents:

$$K^{\pm}(u) = k_1^{\pm}(u + \tfrac{\hbar}{2})^{-1}k_2^{\pm}(u + \tfrac{\hbar}{2}) \qquad H^{\pm}(u) = k_1^{\pm}(u - \frac{\hbar}{2})k_2^{\pm}(u + \frac{\hbar}{2}) \tag{62}$$

$$E(u) = X^+(u + \tfrac{\hbar}{2}) \qquad F(u) = X^-(u + \frac{\hbar}{2}) \tag{63}$$

then all relations among $K^{\pm}(u)$, $H^{\pm}(u)$, $E(u)$ and $F(u)$ can be calculated as follows:

$$[K^{\pm}(u)\ ,\ K^{\pm}(v)] = [K^+(u)\ ,\ K^-(v)] = 0 \tag{64}$$

$$[H^{\pm}(u)\ ,\ H^{\pm}(v)] = [K^{\pm}(u)\ ,\ H^{\pm}(v)] = 0 \tag{65}$$

$$\frac{u_{\mp} - v_{\pm} - \hbar}{u_{\mp} - v_{\pm} + \hbar}H^{\pm}(u)K^{\mp}(v) = K^{\mp}(v)H^{\pm}(u)\frac{u_{\pm} - v_{\mp} - \hbar}{u_{\pm} - v_{\mp} + \hbar} \tag{66}$$

$$\left(\frac{u_{\mp} - v_{\pm} - \hbar}{u_{\mp} - v_{\pm} + \hbar}\right)^2 H^{\pm}(u)H^{\mp}(v) = H^{\mp}(v)H^{\pm}(u)\left(\frac{u_{\pm} - v_{\mp} - \hbar}{u_{\pm} - v_{\mp} + \hbar}\right)^2 \tag{67}$$

$$[K^{\pm}(u)\ ,\ E(v)] = [K^{\pm}(u)\ ,\ F(v)] = 0 \tag{68}$$

$$E(v)H^{\pm}(u) = \frac{u_{\pm} - v + \hbar}{u_{\pm} - v - \hbar}H^{\pm}(u)E(v) \tag{69}$$

$$H^{\pm}(u)F(v) = \frac{u_{\mp} - v + \hbar}{u_{\mp} - v - \hbar}F(v)H^{\pm}(u) \tag{70}$$

$$\{E(u)\ ,\ F(v)\} = \hbar\left[\delta(u_- - v_+)K^+(u_-) - \delta(u_+ - v_-)K^-(v_-)\right] \tag{71}$$

The co-product structure can be derived directly from (6)(7) :

$$\triangle\left(E^{\pm}(u)\right) = E^{\pm}(u) \otimes 1 + H^{\pm}(u \mp \frac{\hbar}{4}c_1) \otimes E^{\pm}(u \mp \frac{\hbar}{2}c_1) \tag{72}$$

$$\triangle\left(F^{\pm}(u)\right) = 1 \otimes F^{\pm}(u) + F^{\pm}(u \pm \frac{\hbar}{2}c_2) \otimes H^{\pm}(u \pm \frac{\hbar}{4}c_2) \tag{73}$$

$$\begin{aligned}\triangle\left(H^{\pm}(u)\right) = {} & H^{\pm}(u \pm \frac{\hbar}{4}c_2) \otimes H^{\pm}(u \mp \frac{\hbar}{4}c_1) - 2F^{\pm}(u \pm \frac{\hbar}{4}c_2 \mp \frac{\hbar}{4}c_1 - \hbar)H^{\pm}(u \pm \frac{\hbar}{4}c_2) \otimes \\ & H(u \mp \frac{\hbar}{4}c_1)^{\pm}E^{\pm}(u \pm \frac{\hbar}{4}c_2 \mp \frac{\hbar}{4}c_1 - \hbar))\end{aligned} \tag{74}$$

$$\triangle\left(K^{\pm}(u)\right) = K^{\pm}(u \pm \frac{\hbar}{4}c_2) \otimes K^{\pm}(u \mp \frac{\hbar}{4}c_1) \tag{75}$$

$$\epsilon(K^{\pm}(u)) = \epsilon(H^{\pm}(u)) = 1 \tag{76}$$

$$\epsilon(E^{\pm}(u)) = \epsilon(F^{\pm}(u)) = 0 \tag{77}$$

The quantum affine superalgebra $U_q(gl(\widehat{1|1}))$ corresponding to the trigonometric solution of super YBE have also been studied in our paper [7] and the general case $U_q(gl(\widehat{m|n}))$ have been independently worked out by Zhang [8].

References

[1] V.G. Drinfel′d, "Quantum Groups", *Proc.ICM-86(Berkeley)*. Vol.1. NewYork Academic Press 1986, 789

[2] V.G. Drinfel′d: *Soviet Math. Dokl.* **36** (1988) 212

[3] N.Yu. Reshetikhin and M.A. Semenov-Tian-Shansky, *Lett. Math. Phys.* **19** (1990) 133

[4] J. Ding and I.B. Frenkel, *Commun. Math. Phys.* **156** (1993) 277

[5] J.F. Cai, G.X. Ju, K. Wu and S.K. Wang, "Super Yangian Double $DY\,(gl(1|1))$ and Its Gauss Decomposition ", preprint, q-alg/**9701011**

[6] L. Liao and X.C. Song, *Mod. Phys. Lett.* **A6** (1991) 959

[7] J.F. Cai, S.K. Wang, K. Wu and W.Z. Zhao, "Drinfel′d Realization of Quantum Affine Superalgebra $U_q(gl\widehat{(1|1)})$", preprint, q-alg/**9703022**

[8] Y.Z. Zhang "Comments on Drinfeld Realization of Quantum Affine Superalgebra $U_q[gl(m|n)^{(1)}]$ and Its Hopf Algebra Structure", preprint, q-alg/**9703020**